





## Boyles' Law

Use Boyle's Law to answer the following questions:

- 1) 1.00 L of a gas at standard temperature and pressure is compressed to 473 mL. What is the new pressure of the gas?
  
  
  
  
  
  
  
  
  
  
- 2) In a thermonuclear device, the pressure of 0.050 liters of gas within the bomb casing reaches  $4.0 \times 10^6$  atm. When the bomb casing is destroyed by the explosion, the gas is released into the atmosphere where it reaches a pressure of 1.00 atm. What is the volume of the gas after the explosion?
  
  
  
  
  
  
  
  
  
  
- 3) Synthetic diamonds can be manufactured at pressures of  $6.00 \times 10^4$  atm. If we took 2.00 liters of gas at 1.00 atm and compressed it to a pressure of  $6.00 \times 10^4$  atm, what would the volume of that gas be?
  
  
  
  
  
  
  
  
  
  
- 4) The highest pressure ever produced in a laboratory setting was about  $2.0 \times 10^6$  atm. If we have a  $1.0 \times 10^{-5}$  liter sample of a gas at that pressure, then release the pressure until it is equal to 0.275 atm, what would the new volume of that gas be?

- 5) Atmospheric pressure on the peak of Mt. Everest can be as low as 150 mm Hg, which is why climbers need to bring oxygen tanks for the last part of the climb. If the climbers carry 10.0 liter tanks with an internal gas pressure of  $3.04 \times 10^4$  mm Hg, what will be the volume of the gas when it is released from the tanks?
- 6) Part of the reason that conventional explosives cause so much damage is that their detonation produces a strong shock wave that can knock things down. While using explosives to knock down a building, the shock wave can be so strong that 12 liters of gas will reach a pressure of  $3.8 \times 10^4$  mm Hg. When the shock wave passes and the gas returns to a pressure of 760 mm Hg, what will the volume of that gas be?
- 7) Submarines need to be extremely strong to withstand the extremely high pressure of water pushing down on them. An experimental research submarine with a volume of 15,000 liters has an internal pressure of 1.2 atm. If the pressure of the ocean breaks the submarine forming a bubble with a pressure of 250 atm pushing on it, how big will that bubble be?
- 8) Divers get “the bends” if they come up too fast because gas in their blood expands, forming bubbles in their blood. If a diver has 0.05 L of gas in his blood under a pressure of 250 atm, then rises instantaneously to a depth where his blood has a pressure of 50.0 atm, what will the volume of gas in his blood be? Do you think this will harm the diver?

## Charles' Law Worksheet

- 1) The temperature inside my refrigerator is about  $4^{\circ}$  Celsius. If I place a balloon in my fridge that initially has a temperature of  $22^{\circ}$  C and a volume of 0.5 liters, what will be the volume of the balloon when it is fully cooled by my refrigerator?
  
- 2) A man heats a balloon in the oven. If the balloon initially has a volume of 0.4 liters and a temperature of  $20^{\circ}$  C, what will the volume of the balloon be after he heats it to a temperature of  $250^{\circ}$  C?
  
- 3) On hot days, you may have noticed that potato chip bags seem to “inflate”, even though they have not been opened. If I have a 250 mL bag at a temperature of  $19^{\circ}$  C, and I leave it in my car which has a temperature of  $60^{\circ}$  C, what will the new volume of the bag be?
  
- 4) A soda bottle is flexible enough that the volume of the bottle can change even without opening it. If you have an empty soda bottle (volume of 2 L) at room temperature ( $25^{\circ}$  C), what will the new volume be if you put it in your freezer ( $-4^{\circ}$  C)?

- 5) Some students believe that teachers are full of hot air. If I inhale 2.2 liters of gas at a temperature of  $18^{\circ}\text{C}$  and it heats to a temperature of  $38^{\circ}\text{C}$  in my lungs, what is the new volume of the gas?
- 6) How hot will a 2.3 L balloon have to get to expand to a volume of 400 L? Assume that the initial temperature of the balloon is  $25^{\circ}\text{C}$ .
- 7) I have made a thermometer which measures temperature by the compressing and expanding of gas in a piston. I have measured that at  $100^{\circ}\text{C}$  the volume of the piston is 20 L. What is the temperature outside if the piston has a volume of 15 L?

## Ideal Gas Law Problems

*Use the ideal gas law to solve the following problems:*

- 1) If I have 4 moles of a gas at a pressure of 5.6 atm and a volume of 12 liters, what is the temperature?
- 2) If I have an unknown quantity of gas at a pressure of 1.2 atm, a volume of 31 liters, and a temperature of 87 °C, how many moles of gas do I have?
- 3) If I contain 3 moles of gas in a container with a volume of 60 liters and at a temperature of 400 K, what is the pressure inside the container?
- 4) If I have 7.7 moles of gas at a pressure of 0.09 atm and at a temperature of 56 °C, what is the volume of the container that the gas is in?
- 5) If I have 17 moles of gas at a temperature of 67 °C, and a volume of 88.89 liters, what is the pressure of the gas?
- 6) If I have an unknown quantity of gas at a pressure of 0.5 atm, a volume of 25 liters, and a temperature of 300 K, how many moles of gas do I have?

- 7) If I have 21 moles of gas held at a pressure of 78 atm and a temperature of 900 K, what is the volume of the gas?
- 8) If I have 1.9 moles of gas held at a pressure of 5 atm and in a container with a volume of 50 liters, what is the temperature of the gas?
- 9) If I have 2.4 moles of gas held at a temperature of 97 °C and in a container with a volume of 45 liters, what is the pressure of the gas?
- 10) If I have an unknown quantity of gas held at a temperature of 1195 K in a container with a volume of 25 liters and a pressure of 560 atm, how many moles of gas do I have?
- 11) If I have 0.275 moles of gas at a temperature of 75 K and a pressure of 1.75 atmospheres, what is the volume of the gas?
- 12) If I have 72 liters of gas held at a pressure of 3.4 atm and a temperature of 225 K, how many moles of gas do I have?